

Vacuum Interpretation for Supergravity M -Branes

Chiang-Mei Chen, Dmitri V. Gal'tsov and Sergei A. Sharakin *

Department of Theoretical Physics, Moscow State University, 119899 Moscow, Russia

A non-local classical duality between the three-block truncated $11D$ supergravity and the $8D$ vacuum gravity with two commuting Killing symmetries is established. The supergravity four-form field is generated via an inverse dualisation of the corresponding Killing two-forms in six dimensions. $11D$ supersymmetry condition is shown to be equivalent to existence of covariantly constant spinors in eight dimensions. Thus any solution to the vacuum Einstein equations in eight dimensions depending on six coordinates and admitting Killing spinors have supersymmetric $11D$ -supergravity counterparts. Using this duality we derive some new brane solutions to $11D$ -supergravity including $1/4$ supersymmetric intersecting M -branes with a NUT parameter and a dyon solution joining the $M2$ and $M5$ -branes intersecting at a point.

To reveal classical symmetries in dimensionally reduced supergravities it is common to use dualisation of the antisymmetric forms involved [1,2]. The standard prescription consists in using this procedure to maximally reduce the rank of forms in a given dimension D , i.e. to dualise those forms whose rank k is greater than $D-k$. Recently it was argued that deeper understanding of hidden symmetries requires more general dualisation schemes, or even introduction of double space of all forms both dualised and non-dualised with the corresponding constraints [3]. Here we want to explore a possibility provided by an *inverse dualisation* ($k < D-k$) to find a non-local correspondence between supergravities involving higher rank antisymmetric forms and vacuum Einstein gravity admitting isometries. The main idea is to dualise the Killing forms associated with these isometries in order to generate higher rank forms which could be then reinterpreted as matter fields of suitable supergravity. For definiteness we consider the correspondence between the three-block truncation of $11D$ -supergravity and $8D$ Einstein gravity with two commuting isometries.

Consider the bosonic sector of $11D$ supergravity

$$S_{11} = \int d^{11}x \sqrt{-\hat{g}_{11}} \left\{ \hat{R}_{11} - \frac{1}{48} \hat{F}_{[4]}^2 \right\} - \frac{1}{6} \int \hat{F}_{[4]} \wedge \hat{F}_{[4]} \wedge \hat{A}_{[3]}, \quad (1)$$

assuming for the space-times the following three-block structure $M^{11} = M^2(z^a) \times M^3(y^i) \times M^{1,5}(x^\mu)$, where M^2 and M^3 are conformally flat euclidean spaces, $M^{1,5}$ is the pseudoriemannian six-dimensional space-time and it is understood that all variables depend only on coordinates x^μ :

$$ds_{11}^2 = g_2^{\frac{1}{2}} \delta_{ab} dz^a dz^b + g_3^{\frac{1}{3}} \delta_{ij} dy^i dy^j + g_2^{-\frac{1}{4}} g_3^{-\frac{1}{4}} g_{\mu\nu} dx^\mu dx^\nu. \quad (2)$$

Here the scalars g_2, g_3 and the six-dimensional metric $g_{\mu\nu}$ are functions of x^μ . The corresponding consistent truncation of $11D$ supergravity involves the following ansatz for the $11D$ four-form $\hat{F}_{[4]} = d\hat{A}_{[3]}$:

$$\begin{aligned} \hat{F}_{[4]\mu\nu ab} &= \epsilon_{ab} F_{\mu\nu}, & \hat{F}_{[4]\mu ijk} &= \epsilon_{ijk} \partial_\mu \kappa, \\ \hat{F}_{[4]\mu\nu\lambda\tau} &= H_{\mu\nu\lambda\tau}, \end{aligned} \quad (3)$$

where $H = dB$ and $F = dA$ are the *six-dimensional* four-form and two-forms respectively.

It is straightforward to show that the equations of motion for the six-dimensional variables can be derived from the following action

$$S_6 = \int \sqrt{-g_6} d^6x \left\{ R_6 - \frac{1}{2} e^{2\phi} (\partial\kappa)^2 - \frac{1}{2} (\partial\phi)^2 - \frac{1}{3} (\partial\psi)^2 - \frac{1}{4} e^{-\phi} \left(e^\psi F^2 + \frac{1}{12} e^{-\psi} H^2 \right) \right\} + \kappa \int F \wedge H, \quad (4)$$

where

$$\phi = -\frac{1}{2} \ln g_3, \quad \psi = -\frac{3}{4} \ln g_2 - \frac{1}{4} \ln g_3. \quad (5)$$

Our main result consists in the following

Theorem: *Any solution of $8D$ vacuum Einstein gravity with two commuting spacelike isometries gives rise to several solutions of the $11D$ supergravity of the form (2,3). If, in addition, the $8D$ space-time admits covariantly constant spinors, the corresponding $11D$ solutions have unbroken supersymmetries.*

To prove these assertions, first we note that the scalar fields κ, ϕ parameterize the coset $SL(2, R)/SO(2)$. To put the action in a manifestly $SL(2, R)$ -covariant form one has to dualise the four-form H as follows:

$$H_{\mu\nu\lambda\tau} = \frac{1}{2} \sqrt{-g} e^{\psi+\phi} \epsilon_{\mu\nu\lambda\tau\rho\sigma} (*H^{\rho\sigma} + \kappa F^{\rho\sigma}). \quad (6)$$

*Email: chen@grg1.phys.msu.su, galtsov@grg.phys.msu.su, sharakin@grg1.phys.msu.su

The equations of motion in the new variables can be derived from the action

$$S_6 = \int \left\{ R_6 - \frac{1}{3}(\partial\psi)^2 - \frac{1}{4}e^\psi \mathcal{F}_{\mu\nu}^T \mathcal{M} \mathcal{F}^{\mu\nu} + \frac{1}{4} \text{Tr}(\partial\mathcal{M}\partial\mathcal{M}^{-1}) \right\} \sqrt{-g_6} d^6x, \quad (7)$$

where

$$\mathcal{M} = \begin{pmatrix} e^{-\phi} + \kappa^2 e^\phi & \kappa e^\phi \\ \kappa e^\phi & e^\phi \end{pmatrix}, \quad \mathcal{F}_{\mu\nu} = \begin{pmatrix} F_{\mu\nu} \\ *H_{\mu\nu} \end{pmatrix}. \quad (8)$$

This action has a manifest $R \times SL(2, R)$ symmetry:

$$\mathcal{M}' = \mathcal{G}^T \mathcal{M} \mathcal{G}, \quad \mathcal{F}'_{\mu\nu} = \mathcal{G}^{-1} \mathcal{F}_{\mu\nu}, \quad \mathcal{G} \in SL(2, R), \quad (9)$$

$$\psi' = \psi + t, \quad \mathcal{F}'_{\mu\nu} = e^{-\frac{t}{2}} \mathcal{F}_{\mu\nu}, \quad t \in R, \quad (10)$$

which can be used to generate new supergravity solutions.

Now let us start with the 8D Einstein theory on space-times with two spacelike Killing symmetries, in which case the 8D metric can be presented as

$$ds_8^2 = h_{ab}(d\zeta^a + \mathcal{A}_\mu^a dx^\mu)(d\zeta^b + \mathcal{A}_\nu^b dx^\nu) + (\det h)^{-\frac{1}{4}} g_{\mu\nu} dx^\mu dx^\nu, \quad (11)$$

where h_{ab} , 2×2 matrix, and $\mathcal{A}_\mu^a$ are two Kaluza-Klein (KK) vectors depending on x^μ only. Under the KK reduction to six dimensions one gets two KK two-forms, and three scalar moduli parameterizing the metric h_{ab} . The eight-dimensional Einstein equations in terms of the six-dimensional variables may be cast exactly into the form (7) provided the following identification holds:

$$h_{ab} = e^{\frac{2}{3}\psi} \begin{pmatrix} e^{-\phi} + \kappa^2 e^\phi & \kappa e^\phi \\ \kappa e^\phi & e^\phi \end{pmatrix}, \quad F = d\mathcal{A}^1, \quad *H = d\mathcal{A}^2. \quad (12)$$

This completes the proof of the bosonic part of the theorem. Note that the four-form $H_{\mu\nu\lambda\tau}$ of the 11D supergravity is generated from the KK two-form by *inverse dualisation*. Possible permutation of two Killing vectors in (11) will lead to different 11D solutions. Moreover, if the actual number of commuting spacelike isometries of the 8D solution is greater than two, one can generate several different 11D solutions via different choice of the ordered pair of ζ^a -directions in (11).

Consider now the 11D supergravity condition for vanishing of the supersymmetry variation of the gravitino field (Killing spinor equation) in a bosonic background satisfying the ansatz (2,3):

$$D_M \epsilon_{11} + \frac{1}{288} (\Gamma_M^{\bar{N}\bar{P}\bar{Q}\bar{R}} - 8\delta_M^{\bar{N}} \Gamma^{\bar{P}\bar{Q}\bar{R}}) \hat{F}_{\bar{N}\bar{P}\bar{Q}\bar{R}} \epsilon_{11} = 0. \quad (13)$$

Here ϵ_{11} is a 32-component Majorana spinor, and the orthonormal components of the gamma-matrices and the four-form are denoted by bar. Consistently with the three-block decomposition for the metric one assumes

the $2 \times 3 \times 6$ split for the spinor $\epsilon_{11} = \epsilon_2 \times \epsilon_3 \times \epsilon_6$ and the 11D gamma-matrices

$$\begin{aligned} \Gamma_{\bar{a}} &= \rho_{\bar{a}} \times \sigma_{\bar{0}} \times \gamma_{\bar{7}}, & \Gamma_{\bar{i}} &= \rho_{\bar{z}} \times \sigma_{\bar{i}} \times \gamma_{\bar{7}}, \\ \Gamma_{\bar{\alpha}} &= \rho_{\bar{0}} \times \sigma_{\bar{0}} \times \gamma_{\bar{\alpha}}, \end{aligned} \quad (14)$$

where $\rho_{\bar{a}}, \sigma_{\bar{i}}$ are two sets of 2×2 Pauli matrices, $a = x, y, i = x, y, z$, $\rho_{\bar{0}}$ and $\sigma_{\bar{0}}$ are unit matrices and $\gamma_{\bar{\alpha}}$ are 6D gamma-matrices: $\{\gamma_{\bar{\alpha}}, \gamma_{\bar{\beta}}\} = 2\eta_{\bar{\alpha}\bar{\beta}}$. The chiral operator is defined as $\gamma_{\bar{7}} = \gamma_{\bar{0}}\gamma_{\bar{1}}\gamma_{\bar{2}}\gamma_{\bar{3}}\gamma_{\bar{4}}\gamma_{\bar{5}}$, so that $\gamma_{\bar{7}}^2 = 1$ and $\{\gamma_{\bar{7}}, \gamma_{\bar{\alpha}}\} = 0$.

Without loss of generality one can assume that ϵ_2 and ϵ_3 are constant spinors. Then after reduction the Eq.(13) is cast into a set of three equations one of which is the 6D differential equation for the spinor ϵ_6 :

$$\left\{ 4D_{\bar{\alpha}} - \frac{1}{3}\partial_{\bar{\alpha}}(\phi + 2\psi) - \frac{1}{3}\gamma_{\bar{\alpha}}^{\bar{\beta}}\partial_{\bar{\beta}}\psi + e^{\frac{\psi+\phi}{2}}\gamma_{\bar{7}}(*H_{\bar{\alpha}} + \kappa F_{\bar{\alpha}}) - is_z \left(e^\phi \gamma_{\bar{7}} \partial_{\bar{\alpha}} \kappa + e^{\frac{\psi-\phi}{2}} F_{\bar{\alpha}} \right) \right\} \epsilon_6 = 0, \quad (15)$$

where $D_{\bar{\alpha}}$ is the six-dimensional spinor covariant derivative and the constant spinor ϵ_2 is split into the chiral projections $\rho_{\bar{z}}\epsilon_2 = s_z\epsilon_2$, while two others are the algebraic equations for ϵ_6 which impose consistency conditions on the bosonic background:

$$\left\{ \not{\partial}(\phi - \frac{2}{3}\psi) - is_z \left(e^\phi \gamma_{\bar{7}} \not{\partial} \kappa + e^{\frac{\psi-\phi}{2}} \mathbf{F} \right) \right\} \epsilon_6 = 0, \quad (16)$$

$$\left\{ \not{\partial}(\phi + \frac{2}{3}\psi) + e^{\frac{\psi+\phi}{2}} \gamma_{\bar{7}}(*\mathbf{H} + \kappa \mathbf{F}) - is_z e^\phi \gamma_{\bar{7}} \not{\partial} \kappa \right\} \epsilon_6 = 0. \quad (17)$$

Here the following notation is used

$$\begin{aligned} \not{\partial} &= \gamma^{\bar{\alpha}} \partial_{\bar{\alpha}} = \gamma^{\bar{\alpha}} e_{\bar{\alpha}}^\mu \partial_\mu, \\ \mathbf{F} &= \frac{1}{2} F_{\bar{\alpha}\bar{\beta}} \gamma^{\bar{\alpha}\bar{\beta}}, \quad F_{\bar{\alpha}} = \frac{1}{2} [\gamma_{\bar{\alpha}}, \mathbf{F}] = F_{\bar{\alpha}\bar{\beta}} \gamma^{\bar{\beta}}, \end{aligned} \quad (18)$$

and similarly for $*H$. In deriving the Eqs. (15-17) all terms containing the products of three, four and five gamma matrices were eliminated via the Fierz identities.

One can show that these equations are equivalent to the geometric Killing spinor equation in $D = 8$:

$$D_M \epsilon_8 = \partial_M \epsilon_8 + \frac{1}{4} \Omega^{\bar{A}\bar{B}}{}_M \mathcal{Y}_{\bar{A}\bar{B}} \epsilon_8 = 0, \quad (19)$$

where ϵ_8 is 16-component spinor, the symbol $\mathcal{Y}_{\bar{A}\bar{B}}$ denotes an antisymmetric product of the 8D gamma-matrices and D_M stands for the spinor covariant derivative with the riemannian connection corresponding to the metric (11). Splitting the 8D spinor as $\epsilon_8 = \epsilon'_2 \times \epsilon'_6$, where ϵ'_2 is a constant spinor, and decomposing the gamma-matrices $\mathcal{Y}_{\bar{a}} = \tau_{\bar{a}} \times \gamma_{\bar{7}}$, $\mathcal{Y}_{\bar{\alpha}} = \tau_{\bar{0}} \times \gamma_{\bar{\alpha}}$, ($\tau_{\bar{a}}$ is the Pauli matrix) one derives from (19) a six-dimensional spinor equation ($M = \mu$) and some matrix constraint equation ($M = a$). The subsequent derivation (details will be given elsewhere) involves an unitary transformation of the corresponding matrix finally arriving at the Eqs.(16-17), and establishing the following correspondence between six-dimensional spinors:

$$\epsilon'_6 = e^{-\frac{\phi+2\psi}{12}} \epsilon_6. \quad (20)$$

This proves the fermionic part of our theorem. Due to the integrability condition for the Eq.(19), the 8D metrics admitting Killing spinors will be Ricci-flat. Also, it is worth noting that the supersymmetry conditions are invariant under symmetry transformations (9).

We do not intend to explore here the general classes of 11D solutions which can be generated via this duality, but rather proceed with some physically meaningful examples. The simplest examples are the non-rotating black $M2$ -brane and $M5$ -brane. Consider the 8D vacuum metric

$$ds_8^2 = H(d\zeta_1 + A_t dt)^2 + d\zeta_2^2 - H^{-1} f dt^2 + dx_1^2 + \cdots + dx_k^2 + f^{-1} dr^2 + r^2 d\Omega_{4-k}, \quad (21)$$

where k is an integer, $(0 \leq k \leq 2)$ and

$$H = 1 + \frac{2\delta}{r^{3-k}}, \quad f = 1 - \frac{2m}{r^{3-k}}, \quad A_t = \frac{2\sqrt{\delta(m+\delta)}}{r^{3-k}H}. \quad (22)$$

Two commuting Killing vector fields may be chosen either as $(\partial_{\zeta_1}, \partial_{\zeta_2})$ or as $(\partial_{\zeta_2}, \partial_{\zeta_1})$. In the first case, performing the steps described above, we arrive at the $M2$ -brane solution

$$ds_{M2}^2 = H^{-\frac{2}{3}} (-f dt^2 + dz_1^2 + dz_2^2) + H^{\frac{1}{3}} (dy_1^2 + dy_2^2 + dy_3^2 + dx_1^2 + \cdots + dx_k^2 + f^{-1} dr^2 + r^2 d\Omega_{4-k}), \quad (23)$$

and $\hat{A}_{tz_1 z_2} = A_t$. In the second case one gets the $M5$ -brane

$$ds_{M5}^2 = H^{-\frac{1}{3}} (-f dt^2 + dz_1^2 + dz_2^2 + dy_1^2 + dy_2^2 + dy_3^2) + H^{\frac{2}{3}} (dx_1^2 + \cdots + dx_k^2 + f^{-1} dr^2 + r^2 d\Omega_{4-k}), \quad (24)$$

with the standard three-form field [8]. For $m = 0$ the 8D metric (21) admits Killing spinors $\epsilon_8 = H^{-\frac{1}{4}} \epsilon_0$ (in cartesian local frame), where ϵ_0 is a constant spinor satisfying $\mathcal{Y}_{\bar{t}\bar{t}} \epsilon_0 = \epsilon_0$. This clearly corresponds to the BPS saturated M -branes. In a similar way one can obtain rotating branes, solutions endowed with waves and KK monopoles as well as some their intersections.

The above construction illustrates that our mapping from 8D to 11D is one to many, *i.e.* an unique 8D vacuum solution can have several 11D supergravity counterparts depending on the choice and the order of two Killing vectors in eight dimensions. This provides a simple way to proliferate 11D solutions: taking one known solution one can find its 8D vacuum partner and then go back to eleven dimensions with a different choice of the Killing vectors involved. For example, two well-known five-dimensional KK solutions, the Brinkmann wave and the KK-monopole, can be combined in a six-dimensional vacuum solution which has six 11D counterparts including all pairwise intersections of four basic solutions: $M2$ -brane, $M5$ -brane, wave and monopole except the $M2 \perp M5$ -brane. One can also find the NUT-generalisations of the rotating M -branes in a way similar to that used in the five-dimensional theory [7].

New interesting solution looking like a non-standard intersection of the $M2$ and $M5$ branes at a point (not over a string as demanded by the usual intersection rules) can be obtained starting with the 5D Kaluza-Klein dyon. The most general dyonic black hole solution of 5D KK theory was found by Gibbons and Wiltshire [4] and its rotating version by Clément [5] and Rasheed [6]. In the static case this is a three-parameter family (with mass, electric and magnetic charges m, q, p):

$$ds_5^2 = \frac{B}{A} (d\zeta_1 + \mathcal{A}_\mu dx^\mu)^2 + d\zeta_2^2 + dx_1^2 + dx_2^2 + \sqrt{\frac{A}{B}} \left\{ \frac{-\Delta}{\sqrt{AB}} dt^2 + \sqrt{AB} \left(\frac{dr^2}{\Delta} + d\Omega_2 \right) \right\}, \quad (25)$$

where $A = r_-^2 - 2dp^2/d_-$, $B = r_+^2 - 2dq^2/d_+$, $r_\pm = r \pm d/\sqrt{3}$, $d_\pm = d \pm \sqrt{3}m$, $\Delta = r^2 - 2mr + p^2 + q^2 - d^2$, and the KK vector field is given by $\mathcal{A}_t = C/B$, $\mathcal{A}_\phi = 2p \cos \theta$, $C = 2qr_-$. The scalar charge d satisfies the cubic equation

$$q^2/d_+ + p^2/d_- = 2d/3. \quad (26)$$

Using the relations (12, 5) and then (2,3), one obtains the 11D solution

$$ds_{11}^2 = \left(\frac{B}{A} \right)^{-\frac{1}{6}} \left\{ \frac{-\Delta}{\sqrt{AB}} dt^2 + \sqrt{AB} \left(\frac{dr^2}{\Delta} + d\Omega_2 \right) \right\} + \left(\frac{B}{A} \right)^{-\frac{2}{3}} (dz_1^2 + dz_2^2) + \left(\frac{B}{A} \right)^{\frac{1}{3}} (dy_1^2 + dy_2^2 + dy_3^2 + dx_1^2 + dx_2^2), \quad (27)$$

$$\hat{A}_{tz_1 z_2} = \mathcal{A}_t, \quad \hat{A}_{\phi z_1 z_2} = \mathcal{A}_\phi.$$

This solution for $p = 0$ reduces to the usual $M2$ -brane while for $q = 0$ — to the $M5$ -brane (localized only on a part of the full transverse space). However, for both q, p non-zero, this is a new dyonic brane: contrary to the Costa $M2 \subset M5$ -brane [8], for which the electric brane lies totally within the magnetic one, and contrary to the intersecting $M2 \perp M5$ -brane (with a common string), now the common part of the world-volumes is zero-dimensional (we suggest the name $M2 \cup M5$). The electric/magnetic charge densities are $Q = 8\pi q$ and $P = 8\pi p$ respectively.

Let us derive the supersymmetry conditions for this solution. As $\kappa = *H = 0$ for it and the Eq.(17) is satisfied identically, the only constraint on the functions A, B, C, Δ is due to the Eq.(16) which may be represented as

$$\left\{ I_6 + i s_z \left[\alpha(r) \gamma^{\bar{t}} - \beta(r) \gamma^{\bar{r}\bar{\theta}\bar{\varphi}} \right] \right\} \epsilon_6 = 0, \quad (28)$$

where

$$\alpha(r) = \frac{B^2 \partial_r (C/B)}{A^{3/2} \Delta^{1/2} \partial_r (B/A)}, \quad \beta(r) = \frac{2p B^{3/2}}{A^2 \Delta^{1/2} \partial_r (B/A)}.$$

Non-zero solution to (28) can exist if only $\alpha^2(r) + \beta^2(r) = 1$, what implies asymptotically

$$p^2 + q^2 = 4d^2/3. \quad (29)$$

Combining this with the cubic constraint (26) we find that either i) $p = 0$, $m = d/\sqrt{3}$, $q = 2d/\sqrt{3}$, corresponding to an extremal $M2$ -brane (23), or ii) $q = 0$, $m = -d/\sqrt{3}$, $p = 2d/\sqrt{3}$, an extremal $M5$ -brane (24), otherwise the constraints lead to pure vacuum. So the only nontrivial supersymmetric members of the family of composite $M2 \cup M5$ -branes are the single extremal $M2$ and $M5$ -branes.

One can also obtain the rotating version of $M2 \cup M5$ -brane (with one rotational parameter) through the following transformation:

$$\begin{aligned} dt &\rightarrow dt + \omega d\phi, \quad \Delta \rightarrow \Delta + a^2, \\ A &\rightarrow A + a^2 \cos^2 \theta + \frac{2jpq \cos \theta}{d_+^2/3 - q^2}, \\ B &\rightarrow B + a^2 \cos^2 \theta - \frac{2jpq \cos \theta}{d_-^2/3 - p^2}, \\ C &\rightarrow C - \frac{2jp(d_+/\sqrt{3}) \cos \theta}{d_-^2/3 - p^2}, \end{aligned} \quad (30)$$

where

$$\omega = \frac{2j \sin^2 \theta}{\Delta - a^2 \sin^2 \theta} \left[r - m + \frac{d_+(m^2 + d^2 - p^2 - q^2)}{\sqrt{3}(d_+^2/3 - q^2)} \right].$$

In terms of new variables one component of the KK vector potential, $\mathcal{A}_t$, remains the same while another becomes

$$\begin{aligned} \mathcal{A}_\phi &= \frac{C}{B} \omega + \frac{2p\Delta \cos \theta}{\Delta - a^2 \sin^2 \theta} \\ &+ \frac{2jq \sin^2 \theta [(rd_- - md_-)/\sqrt{3} + p^2 + q^2 - d^2]}{(\Delta - a^2 \sin^2 \theta)(d_+^2/3 - q^2)}. \end{aligned} \quad (31)$$

The angular momentum of the system, j , is given by

$$j^2 = a^2 \frac{(d_+^2/3 - q^2)(d_-^2/3 - p^2)}{m^2 + d^2 - p^2 - q^2}. \quad (32)$$

The drawback of our generating technique is that the $11D$ branes obtained can not *localize* on all coordinates of transverse space, since our basic ansatz (2) is not general enough. However one still can use it to explore possible geometric $11D$ structures and then to try to localize solutions in a straightforward way. Here we present without derivation (details will be given in a forthcoming paper) a new partially localized intersecting $M2$ branes with a NUT parameter:

$$\begin{aligned} d\hat{s}_{11}^2 &= (F_1 F_2)^{\frac{1}{3}} \left\{ -(F_1 F_2)^{-1} (dt - \omega_i dx^i)^2 + F_1^{-1} (dz_1^2 + dz_2^2) \right. \\ &\quad \left. + F_2^{-1} (dw_1^2 + dw_2^2) + dy_1^2 + dy_2^2 + dy_3^2 + \delta_{ij} dx^i dx^j \right\}, \\ \hat{A}_{[3]} &= F_1^{-1} (dt - \omega_i dx^i) \wedge dz_1 \wedge dz_2 + U dy_1 \wedge dy_2 \wedge dy_3 \\ &\quad + F_2^{-1} (dt - \omega_i dx^i) \wedge dw_1 \wedge dw_2, \end{aligned} \quad (33)$$

where $U(x)$, $F_2(x)$, are the harmonic functions on the x^i space while $\omega_i(x)$ and $F_1(x, w)$ satisfy the equations

$$\epsilon_{ijk} \partial_j \omega_k = \partial_i U, \quad \partial_x^2 F_1 + F_2 \partial_w^2 F_1 = 0. \quad (34)$$

This solution (contrary to its version without NUT) involves the Chern-Simons term of (1). Direct check reveals the existence of the $8D$ covariantly constant spinors corresponding to the $1/4$ unbroken supersymmetry.

Thus we have given a purely vacuum interpretation to a certain class of the $11D$ supergravity M -branes via new non-local duality between $11D$ supergravity and $8D$ vacuum Einstein gravity. It is worth noting that the very natural correspondence between the Killing spinor equations in both theories may have a deeper meaning in the context of $8D$ supergravities. In this connection it has to be emphasized that the type of duality described here is by no means related to the direct dimensional reduction from $11D$ to $8D$ theory.

One of the authors (DVG) would like to thank the Laboratory of Gravitation and Cosmology of the University Paris-6 for hospitality and the CNRS for support while this work was in progress.

-
- [1] B. Julia, *Group Disintegrations*, in: *Superspace and Supergravity*, Eds. S.W. Hawking and M. Rocek (Cambridge Univ. Press, 1981) 331; *Dualities in the Classical Supergravity Limits*, [hep-th/9805083](#).
 - [2] K.S. Stelle, *Lectures on Supergravity p -Branes*, [hep-th/9710119](#).
 - [3] E. Cremmer, B. Julia, H. Lü and C.N. Pope, *Dualisation of Dualities, I*, *Nucl. Phys.* **B523** (1998) 73-144; [hep-th/9710119](#); *Dualisation of Dualities, II: Twisted Self-Duality of Doubled Fields and Superdualities*, *Nucl. Phys.* **B535** (1998) 242-292; [hep-th/9806106](#).
 - [4] G. Gibbons and D. Wiltshire, *Black Holes in Kaluza-Klein Theory*, *Ann. Phys.* **167** (1986) 201-223; Erratum: *ibid* **176** (1987) 393.
 - [5] G. Clément, *Rotating Kaluza-Klein Monopoles and Dyons*, *Phys. Lett.* **A118** (1986) 11-13; *Stationary Solutions in Five-Dimensional General Relativity*, *Gen. Rel. Grav.* **18** (1986) 137-160.
 - [6] D. Rasheed, *The Rotating Dyonic Black Holes of Kaluza-Klein Theory* *Nucl. Phys.* **B454** (1995) 379-401; [hep-th/9505038](#).
 - [7] C.-M. Chen, D.V. Gal'tsov, K. Maeda and S.A. Sharakin, *$SL(4, R)$ Generating Symmetry in Five-Dimensional Gravity Coupled to Dilaton and Three-Form*, *Phys. Lett.* **B453** (1999) 7-16; [hep-th/9901130](#).
 - [8] M.S. Costa, *Composite M -branes*, *Nucl. Phys.* **B490** (1997) 202-216; [hep-th/9609181](#); *Black Composite M -branes*, *Nucl. Phys.* **B495** (1997) 195-205; [hep-th/9610138](#).